

The Occurrence of Varicella After Zoster Vaccination in a Patient With Rheumatoid Arthritis Receiving Tofacitinib

Chung-Mao KAO , Tsu-Yi HSIEH , Kuo-Lung LAI 

Department of Internal Medicine, Division of Allergy, Immunology and Rheumatology, Taichung Veterans General Hospital, Taichung City, Taiwan

Tofacitinib is a Janus kinase (JAK) 1 and 3 inhibitor approved for the treatment of rheumatoid arthritis (RA).¹ JAK1 is associated with the receptors for interferons (IFNs) which are involved in the immunity against viral infections.^{2,3} Tofacitinib modulates the varicella-zoster virus (VZV)-specific type 1 helper T cells' response to VZV by reducing IFN- γ production and the proliferation of specific T cells in a dose-dependent manner.^{4,5} Common adverse events of tofacitinib include herpes zoster.⁶ Zoster vaccine is one type of live-attenuated vaccines, of which the administration concurrent with the use of tofacitinib is suggested to be avoided.⁷

A 30-year-old female patient diagnosed with RA was administered with tofacitinib 5 mg twice daily. She did not have history of varicella at a young age. Due to personal reasons, she received a single dose of zoster vaccine without consultation with her rheumatologist. Two weeks after the vaccination, itchy vesicles and papules originated from the perioral region and around the vaccine injection site in the left deltoid region (Figure 1). Subsequently, the vesicles and papules rapidly extended to the forehead, cheeks and chin (Figure 2). Varicella in association with

zoster vaccination was considered, so tofacitinib was discontinued. She was admitted to the isolation ward and given intravenous acyclovir 500 mg every 8 h. On hospital Day 1, the results of blood survey were as follows: white blood cell 7,180/ μ L (neutrophils 90.1%, lymphocytes 3.5%), hemoglobin 11.3 g/dL, platelet count 251,000/ μ L, aspartate aminotransferase 56 U/L, alanine aminotransferase 61 U/L, blood urea nitrogen 10 mg/dL, and creatinine



Figure 1. Several vesicles and papules occurred in a group around injection site of zoster vaccine in left deltoid region. Lesions had been crusted when photograph was taken.

Received: June 22, 2020 **Accepted:** July 16, 2020 **Published online:** January 14, 2021

Correspondence: Kuo-Lung Lai, MD, Department of Internal Medicine, Division of Allergy, Immunology and Rheumatology, Taichung Veterans General Hospital, 40705 Taichung City, Taiwan. Tel: +886-4-2359-2525 / 3354 e-mail: kllaichiayi@yahoo.com.tw

Citation:

Kao CM, Hsieh TY, Lai KL. The Occurrence of Varicella After Zoster Vaccination in A Patient With Rheumatoid Arthritis Receiving Tofacitinib. Arch Rheumatol 2021;36(x):i-iii.

©2020 Turkish League Against Rheumatism. All rights reserved.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes (<http://creativecommons.org/licenses/by-nc/4.0/>).



Figure 2. Multiple vesicles and papules occurred over forehead, cheeks, chin and neck. Some lesions had become crusted.

0.74 mg/dL. A serological survey revealed an elevation of VZV-immunoglobulin (Ig) M (ratio=3.887, reference positive ≥ 1.1) and VZV-IgG (ratio=6.366, reference positive ≥ 1.1) levels, along with normal herpes simplex virus (HSV)-I IgG and HSV-II IgG levels. During the first three days of hospitalization, the vesicles continued to extend to the skin covering the neck, upper chest and upper back, which was accompanied by a fever. The fever later subsided and the vesicles stopped their progression and began to become crusted during hospital Day 3. After a seven-day course of intravenous acyclovir therapy, the patient was allowed to be discharged from the isolation ward. No VZV reactivation was observed during the subsequent six-month outpatient department visits. A written informed consent was obtained from the patient.

To our knowledge, this is the first case in the English literature which demonstrates the risk of post-zoster vaccination varicella in a tofacitinib user. The 2019 European League Against Rheumatism recommendations stated that live-attenuated vaccines may be considered

in patients with rheumatic diseases but they should be avoided during immunosuppression.⁷ Zoster vaccination with caution in appropriately-selected patients with rheumatic diseases can be both immunogenic and safe.⁸ For the purpose of safety, it is suggested that the zoster vaccine should be administered four weeks prior to the initiation of biologic or targeted synthetic disease-modifying antirheumatic drugs.⁷ Our case has demonstrated an adverse event of zoster vaccination during tofacitinib treatment. The disease course of zoster vaccine-induced varicella seemed benign and the patient displayed a good response to intravenous acyclovir treatment.

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Hodge JA, Kawabata TT, Krishnaswami S, Clark JD, Telliez JB, Dowty ME, et al. The mechanism of action of tofacitinib - an oral Janus kinase inhibitor for the treatment of rheumatoid arthritis. *Clin Exp Rheumatol* 2016;34:318-28.
2. Pesu M, Laurence A, Kishore N, Zwillich SH, Chan G, O'Shea JJ. Therapeutic targeting of Janus kinases. *Immunol Rev* 2008;223:132-42.
3. O'Sullivan LA, Liongue C, Lewis RS, Stephenson SE, Ward AC. Cytokine receptor signaling through the Jak-Stat-Socs pathway in disease. *Mol Immunol* 2007;44:2497-506.
4. Asanuma H, Sharp M, Maecker HT, Maino VC, Arvin AM. Frequencies of memory T cells specific for varicella-zoster virus, herpes simplex virus, and cytomegalovirus by intracellular detection of cytokine expression. *J Infect Dis* 2000;181:859-66.
5. Almanzar G, Kienle F, Schmalzing M, Maas A, Tony HP, Prelog M. Tofacitinib modulates the VZV-specific CD4+ T cell immune response in vitro in lymphocytes of patients with rheumatoid arthritis. *Rheumatology (Oxford)* 2019;58:2051-60.
6. Pfizer Inc.: Xeljanz prescribing information. Available at: <http://labeling.pfizer.com/ShowLabeling.aspx?id=959>. [Last revised: December 2019].

7. Furer V, Rondaan C, Heijstek MW, Agmon-Levin N, van Assen S, Bijl M, et al. 2019 update of EULAR recommendations for vaccination in adult patients with autoimmune inflammatory rheumatic diseases. *Ann Rheum Dis* 2020;79:39-52.
8. Winthrop KL, Wouters AG, Choy EH, Soma K, Hodge JA, Nduaka CI, et al. The safety and immunogenicity of live zoster vaccination in patients with rheumatoid arthritis before starting tofacitinib: A randomized phase II trial. *Arthritis Rheumatol* 2017;69:1969-77.