

Genetic Polymorphism of Manganese Superoxide Dismutase in Behçet's Disease

Efkan UZ,¹ H. Ramazan YILMAZ,² Ramazan YAĞCI,³ İsmail AKYOL,⁴ Tuğba ERSOY,⁵
Gülten SUNGUR,⁵ Ayşe YİĞİT,¹ Sunay DUMAN,⁵ Ömer AKYOL⁶

¹Department of Medical Genetics, Medical Faculty of Süleyman Demirel University, Isparta, Turkey

²Department of Medical Genetics, Medical Faculty of Mevlana University, Konya, Turkey

³Department of Ophthalmology, Medical Faculty of Pamukkale University, Denizli, Turkey

⁴Animal Science Genetics Unit, Kahramanmaraş Sütçü İmam University Faculty of Agriculture, Kahramanmaraş, Turkey

⁵Department of Ophthalmology, Ankara Training and Research Hospital, Ankara, Turkey

⁶Department of Medical Biochemistry, Medical Faculty of Hacettepe University, Ankara, Turkey

ABSTRACT

Objectives: This study aims to investigate the genetic association between single nucleotide mutation in mitochondrial manganese superoxide dismutase and a Behçet's disease (BD) population by using molecular techniques.

Patients and methods: Ninety-three BD patients (45 males, 48 females; mean age 33.15±8.99 years; range 17 to 65 years) and 125 controls (58 males, 67 females; mean age 28.33±7.31 years; range 18 to 62 years) were genotyped by polymerase chain reaction-restriction fragment length polymorphism method. The genotypic distributions in BD patients and controls were consistent with the Hardy-Weinberg equilibrium.

Results: Significant differences were observed between BD patients and controls in terms of genotypic distribution. Frequencies of alanine (Ala)/Ala, Ala/valine (Val), and Val/Val were 14.0% (n=13), 45.2% (n=42), and 40.9% (n=38) in BD patients and 21.6% (n=27), 53.6% (n=67), and 24.8% (n=31) in controls, respectively (p=0.033).

Conclusion: The Val/Val genotype of the manganese superoxide dismutase gene is associated with the physiopathology of BD in a group of Turkish patients.

Keywords: Behçet's disease; manganese superoxide dismutase; polymerase chain reaction-restriction fragment length polymorphism; polymorphism.

Behçet's disease (BD) is a chronic, multi-systemic, relapsing, and inflammatory disorder characterized by ocular manifestations, in particular.¹ It is also characterized by uveitis, recurrent oral ulceration, genital ulceration, and skin lesions. BD exists worldwide, but it is relatively more widespread in Middle Eastern and East Asian countries including Japan.² The effect of autoimmune response and genetic factors on etiology of BD was well established.² However, the etiologic mechanisms underlying this vascular disease have not been well understood yet. Reactive oxygen species (ROS) have been hypothesized to play

an important role in BD. Evidence from several studies suggests that neutrophils from patients with BD generate high levels of ROS and then this stress causes endothelial cell lysis *in vitro*.³⁻⁵ On the other hand, different data reported in the literature show a variation for the activities of antioxidant enzymes in BD: erythrocyte superoxide dismutase (SOD) activity was found to be increased,^{4,6,7} unchanged,⁸ and decreased,^{5,9} glutathione reductase activity was found to be increased,⁷ catalase, glutathione reductase and glutathione peroxidase (GPx) activities were found to be decreased,⁵⁻⁹ and ultimately, erythrocyte

Received: April 20, 2015 **Accepted:** August 09, 2015 **Published online:** December 12, 2015

Correspondence: Efkan Uz, MD. Süleyman Demirel Üniversitesi Tıp Fakültesi Tıbbi Genetik Anabilim Dalı, 32260 Isparta, Turkey.

Tel: +90 246 - 211 32 34 e-mail: efkanuz@yahoo.com

©2016 Turkish League Against Rheumatism. All rights reserved.

catalase activity was found to be unchanged⁴ in BD patients.

Superoxide dismutase is an enzyme catalyzing the dismutation reaction of superoxide radicals to hydrogen peroxide.^{10,11} SOD, a key antioxidant enzyme, is a family of enzymes, comprising copper, zinc-SOD, manganese (Mn)-SOD, and extracellular SOD, whose function is protection against ROS.^{2,12} Mn-SOD gene is located on chromosome 6q25 in mitochondria and it consists of five exons. Valine-to-alanine (Val-to-Ala) substitution at codon 16 of human Mn-SOD might lead to misdirected intracellular trafficking followed by changes in Mn-SOD activity in the mitochondria.^{2,13,14} In this study, we aimed to investigate the genetic association between single nucleotide mutation in mitochondrial Mn-SOD and a BD population by using molecular techniques.

PATIENTS AND METHODS

This multicenter study included 93 BD patients (45 males, 48 females; mean age 33.15±8.99 years; range 17 to 65 years) and 125 controls (58 males, 67 females; mean age 28.33±7.31 years; range 18 to 62 years) between October 2004 and July 2005. Clinical and demographic data were recorded. The study was approved by the local non-invasive clinical research ethics committee. All of the participants provided written informed consent. Patients were diagnosed according to the International Study Group for BD.¹⁵

Blood samples (5-10 mL), taken in ethylenediaminetetraacetic acid containing tubes during vascular access established for the normal clinical biochemistry tests of participants, were obtained from volunteer BD patients. Characteristics of patient and control groups were summarized in Table 1. Genomic deoxyribonucleic

acid (DNA) was isolated from 300 µL blood using Promega Wizard Genomic DNA purification Kit (Promega Corp., Madison, WI, USA). Isolation process was performed according to the manufacturer's recommendations.¹⁶ The DNA was dissolved in 1X TE buffer by mixing and stored at -20 °C.

The signal sequence-containing region was amplified by polymerase chain reaction (PCR) using Mn-SOD forward and reverse primers in accordance with previous reports.¹⁶⁻¹⁸ Oligonucleotides were purchased from Iontek (Iontek Medical A.Ş. Moleküler Biyoloji ve Genetik Laboratuvarı, Istanbul, Turkey). An Ala/Val polymorphism in the signal peptide of Mn-SOD gene was evaluated using a primer pair (forward 5' ACCAGCAGGCAGCTGGCGCCGG 3' and reverse 5' GCGTTGATGTGAGGT TCCAG 3') to amplify a 107 bp fragment followed by digestion with the NgoM IV gene from *Neisseria gonorrhoeae* MS11 (NgoM IV) recognizes the nucleotide sequence as a restriction site GCCGGC (Figure 1). PCR amplification of genes was performed in a total volume of 50 µL, containing 50 ng of genomic DNA, 20 pmol µL⁻¹ of each primer, 1.25 U Taq polymerase (in 50 mM Tris-hydrochloride, 100 mM sodium chloride, 0.1 mM ethylenediaminetetraacetic acid, 1 mM dithiothreitol, 50% glycerol and 1% Triton X-100), 2 mM deoxy nucleotide triphosphate, 2 mM MgCl₂, 1X PCR buffer containing 50 mM potassium chloride, 10 mM Tris-hydrochloride (pH 8.3 at 25 °C) (Promega Madison, WI, USA). The PCR reaction conditions involved an initial denaturation of DNA at 95 °C for five minutes, followed by 35 cycles of amplification at 95 °C for one minute (melting), 61 °C for one minute (annealing) and 72 °C for two minutes, and final extension at 72 °C for seven minutes in a thermal cycler. The resulting 107 bp PCR product

Table 1. Main characteristics of collected samples, genotypes of Ala-9Val polymorphism and allele frequencies in Behçet's disease patients and controls

	Sample descriptions			Genotypes*						Allele frequencies**	
	n	Gender		Ala/Ala		Ala/Val		Val/Val		Ala	Val
		Male/Female		n	%	n	%	n	%	p	p
Behçet's disease	93	45/48		13	14.0	42	45.2	38	40.9	0.366	0.634
Control	125	58/67		27	21.6	67	53.6	31	24.8	0.484	0.516

n: Number of collected or observed samples; * Significant difference in genotype frequencies between patients and controls ($\chi^2=6.793$, $df=2$, $p=0.033$).
** No significant difference in allele frequencies between patients and controls.

was digested with the restriction endonuclease NgoM IV at 37 °C for 16 hours according to the manufacturer's recommendations (5 µL PCR product was digested in total volume of 10 µL using buffer including bovine serum albumin and 1 µL NgoM IV (5 U µL⁻¹) restriction enzyme.

Isolated chromosomal DNAs were run on 1% agarose gel at 100V for an hour and stained with ethidium bromide (0.5 g µL⁻¹). Tris-Borate-ethylenediaminetetraacetic acid buffer (0.5X) was used as running buffer. Digested PCR product was run on 3% MetaPhor agarose. Restriction enzyme digestion results in a 107 bp product (allele 1 Val-9) or 89 and 18 bp products (allele 2 Ala-9).

Statistical analysis

Allele and genotype frequencies between BD patients and controls were compared by Chi-square test using the Statistical Package for the Social Sciences (SPSS) for Windows version 16.0

software program (SPSS Inc., Chicago, IL, USA). Allele frequencies were calculated by Internet based Pop-Gen 3.2 software. Differences in the demographic characteristics of patients were assessed among the three Mn-SOD genotypes using one-way analyses of variance (ANOVA). The Chi-square test was used to compare the distribution of genotypes and allele frequencies for deviations from Hardy-Weinberg equilibrium.

RESULTS

Analysis results showed that maximum cleavage site was between position 18 and 19 (Figure 1).

When restriction amplification region had band sizes of 89 bp and 18 bp, this genotype was called homozygote (Val/Val). When restriction resulted in band sizes of 107 bp, 89 bp, and 18 bp, this




Figure 1. Schematic representation of the 661 nucleotide base pairs and its amino acid sequence alignment of human manganese superoxide dismutase gene. Square brackets represent signal sequence. Italic nucleotides represent forward and reverse sequences, respectively. Framed sequence represents the definition sequence of restriction endonuclease NgoM IV on the polymerase chain reaction product. If the last nucleotide of this sequence T converts C (C/C), then NgoM IV can definite this point and cut. In this case, the last product of digestion by NgoM IV is two DNA pieces, 89 and 18 bp (GCT codon codes Ala amino acid). If the last nucleotide remains as T (T/T), NgoM IV cannot definite the sequence and this 107 bp DNA sequence remains intact (GTT codon codes Val amino acid). Asterix (*) represents polymorphic nucleotide. Up-down vertical arrow (↓) represents location of restriction site by NgoM IV.

genotype was called heterozygote (Ala/Val). The allele and genotype frequencies of the subjects for the Ala-9Val Mn-SOD polymorphism were shown in Table 1. If the -9 codon was GCT (Ala), then digestion with NgoM IV produced two DNA fragments, 89 and 18 bp in length. If codon was GTT (Val), the amplified product was not digested with NgoM IV and remained as a whole 107 bp DNA fragment. Genotype of Ala-9Val polymorphism in BD and control groups were shown in Table 1. The genotype distributions of BD and control groups were within the Hardy-Weinberg equilibrium. There was no sex or age-dependent difference between the groups. Significant difference was found in genotype frequencies between BD and control groups ($X^2=6.793$, $df=2$, $p=0.033$). The frequency of Val-9 was more common both in BD patients and controls. No significant difference was found in allele frequencies between BD patients and controls (Table 1, Figure 2).

DISCUSSION

Proteins secreted out of the cell generally contain an intrinsic signal that governs their passage across membranes and cell walls. Specific amino acid sequences determine whether a protein will




Figure 2. Polymerase chain reaction-restriction fragment length polymorphism analysis of Ala-9Val polymorphism in the manganese superoxide dismutase gene. Lanes 1-4 represent four samples after digestion with NgoM IV. line 2: Ala/Ala genotype, line 3: Val/Val genotype, line 4: Ala/Val genotype. M: deoxyribonucleic acid marker, 200 and 100 bp bands represent the related part of deoxyribonucleic acid marker. Bands represent 107 bp (up) and 89 bp (down) deoxyribonucleic acid fragments in lanes 1-4. Last bands (18 bp) in digested polymorphism chain reaction products by NgoM IV are not shown because it is in the bottom of gel together with surplus primers.

pass through a membrane, become integrated into the membrane, or be exported out of the cell. These signal sequences are present either as a short tail at one end of the protein or sometimes located within the protein. The method program is available under the prediction server page of Center for Biological Sequence analysis (<http://www.cbs.dtu.dk/services/SignalP/>). SignalP looks at charge distribution and a unique composition of the hydrophobic region.

It is known that Mn-SOD gene is polymorphic.¹³ We identified a significant association between development of BD disease and Mn-SOD Val-9Ala polymorphism. The Val/Val genotype of the Mn-SOD gene can be a novel genetic marker of risk factor for BD in Turkish people. Our results were similar to those of Nakao et al.,² who showed a significant difference in the allele and genotype frequencies of the Mn-SOD Ala16Val polymorphism between BD patients and control subjects in Japan. On the other hand, Yen et al.¹⁹ found no significant difference in the frequencies of Mn-SOD gene polymorphisms between BD patients and controls in Taiwan.

Many studies were carried out on Mn-SOD polymorphisms in various diseases. While some of these studies report a significant similarity between polymorphisms in patients and controls, some does not. Cox et al.²⁰ reported that Mn-SOD gene was associated with an increased risk of breast cancer. Whereas Lightfoot et al.²¹ suggested that no association was observed between Val16Ala and total non-Hodgkin's lymphoma, diffuse large-B cell lymphoma or follicular lymphoma in the UK and USA. Akyol et al.¹⁸ suggested that when the patients with schizophrenia were divided into three subgroups as disorganized, paranoid and residual, there was a significant difference in Mn-SOD genotypic distribution among the subgroups. It has been shown that SOD Ala-16Val polymorphism is an age-dependent modulator of oxidized low-density lipoprotein cholesterol levels in middle-aged males and elderly females.²² The Ala-9Val polymorphism in the mitochondrial targeting sequence of the Mn-SOD gene is not associated with juvenile-onset asthma in Turkish population.²³ The Ala-9Val polymorphisms in the mitochondrial targeting sequence may influence the efficiency of Mn-SOD transport into mitochondria.¹⁸ It has been shown that Mn-SOD genotypes containing the variant A allele were

found to be associated with a 1.5-fold increased risk of breast cancer compared with those with the homozygous wild-type genotype (Mn-SOD VV).¹⁷

Taysi et al.⁷ found that erythrocyte SOD activities were increased in BD compared to healthy controls whereas the other antioxidant enzymes, glutathione S transferase, catalase, and GPx were decreased. In another study, which investigated the oxidant-antioxidant status in whole blood from BD and controls, GPx was found to be decreased similarly to the previous paper.⁸ Aydin et al.²⁴ found increased malondialdehyde levels, decreased GPx activity and non-significant changes in SOD activity in the serum of the patients. SOD activity was demonstrated to be increased in erythrocyte from BD patients in another study.⁴ All of the above-mentioned studies suggest that there is a changing pattern of oxidative stress and antioxidant system in BD and antioxidant system in plasma, erythrocyte, and other body fluids and cells have been commonly declined. This is important because there is huge amount of oxidative stress parameters which were produced by different metabolic processes in physiological metabolism of the body that need to be scavenged by SOD and other antioxidant enzymes.²⁵⁻²⁷ Therefore, we aimed to show the genetic clues and supporting data on this topic.

Nakao et al.² investigated the association of Mn-SOD and extracellular SOD gene polymorphism as well as endothelial and inducible nitric oxide synthase genes with susceptibility to BD in Japan. They found increased Mn-SOD Val16 frequencies in patients with BD. They alleged that Mn-SOD V/V genotype and human leukocyte antigen-B*5101 had a synergistic role in controlling susceptibility to BD. There were no significant differences in the frequencies of extracellular SOD gene polymorphism (Arg213Gly) as well as nitric oxide synthase gene polymorphism between patients and controls.² Our results conform with results of this study and give us some important clues about the underlying mechanism of BD in relation to oxidative stress. Another research on the same topic from Taiwan reported no significant difference in the allele and genotype frequencies of the Mn-SOD Ala/Val polymorphism between BD and control.¹⁹

These differences between the studies might be originated from the racial and ethnic differences for the related genes.

Valine to alanine substitution may increase the targeting efficiency by a conformational change of the targeting sequence, consequently leading to increased mitochondrial ROS scavenging capability.²⁸ The presence of more than one signal sequence for this enzyme suggests a combinatorial mechanism determining the rates of targeting, membrane translocation and/or signal sequence cleavage with concomitant folding of Mn-SOD.¹³ Sutton et al.²⁹ showed that the Ala-Mn-SOD/mitochondrial targeting sequence allows efficient Mn-SOD import into the mitochondrial matrix, while the Val-variant causes partial arrest of the precursor within the inner membrane and decreased formation of the active Mn-SOD tetramer in the mitochondrial matrix. In conclusion, the results of this study suggest that the Val/Val genotype of the Mn-SOD gene is associated with the physiopathology of BD in a group of Turkish patients.

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Sögüt S, Aydin E, Elyas H, Aksoy N, Ozyurt H, Totan Y, et al. The activities of serum adenosine deaminase and xanthine oxidase enzymes in Behçet's disease. *Clin Chim Acta* 2002;325:133-8.
2. Nakao K, Isashiki Y, Sonoda S, Uchino E, Shimonagano Y, Sakamoto T. Nitric oxide synthase and superoxide dismutase gene polymorphisms in Behçet disease. *Arch Ophthalmol* 2007;125:246-51.
3. Niwa Y, Miyake S, Sakane T, Shingu M, Yokoyama M. Auto-oxidative damage in Behçet's disease-endothelial cell damage following the elevated oxygen radicals generated by stimulated neutrophils. *Clin Exp Immunol* 1982;49:247-55.
4. Gunduz K, Ozturk G, Sozmen EY. Erythrocyte superoxide dismutase, catalase activities and plasma nitrite and nitrate levels in patients with Behçet disease and recurrent aphthous stomatitis. *Clin Exp Dermatol* 2004;29:176-9.

5. Erkiçiç K, Evreklioglu C, Cekmen M, Ozkiris A, Duygulu F, Dogan H. Adenosine deaminase enzyme activity is increased and negatively correlates with catalase, superoxide dismutase and glutathione peroxidase in patients with Behçet's disease: original contributions/clinical and laboratory investigations. *Mediators Inflamm* 2003;12:107-16.
6. Dincer Y, Alademir Z, Hamuryudan V, Fresko I, Akcay T. Superoxide dismutase activity and glutathione system in erythrocytes of men with Behçet's disease. *Tohoku J Exp Med* 2002;198:191-5.
7. Taysi S, Demircan B, Akdeniz N, Atasoy M, Sari RA. Oxidant/antioxidant status in men with Behçet's disease. *Clin Rheumatol* 2007;26:418-22.
8. Buldanlioglu S, Turkmen S, Ayabakan HB, Yenice N, Vardar M, Dogan S, et al. Nitric oxide, lipid peroxidation and antioxidant defence system in patients with active or inactive Behçet's disease. *Br J Dermatol* 2005;153:526-30.
9. Saglam K, Serce AF, Yilmaz MI, Bulucu F, Aydin A, Akay C, et al. Trace elements and antioxidant enzymes in Behçet's disease. *Rheumatol Int* 2002;22:93-6.
10. Akyol S, Erdogan S, Idiz N, Celik S, Kaya M, Ucar F, et al. The role of reactive oxygen species and oxidative stress in carbon monoxide toxicity: an in-depth analysis. *Redox Rep* 2014;19:180-9.
11. Ucar F, Sezer S, Erdogan S, Akyol S, Armutcu F, Akyol O. The relationship between oxidative stress and nonalcoholic fatty liver disease: Its effects on the development of nonalcoholic steatohepatitis. *Redox Rep* 2013;18:127-33.
12. Wang LM, Kitteringham N, Mineshita S, Wang JZ, Nomura Y, Koike Y, et al. The demonstration of serum interleukin-8 and superoxide dismutase in Adamantiades-Behçet's disease. *Arch Dermatol Res* 1997;289:444-7.
13. Rosenblum JS, Gilula NB, Lerner RA. On signal sequence polymorphisms and diseases of distribution. *Proc Natl Acad Sci U S A* 1996;93:4471-3.
14. Shimoda-Matsubayashi S, Matsumine H, Kobayashi T, Nakagawa-Hattori Y, Shimizu Y, Mizuno Y. Structural dimorphism in the mitochondrial targeting sequence in the human manganese superoxide dismutase gene. A predictive evidence for conformational change to influence mitochondrial transport and a study of allelic association in Parkinson's disease. *Biochem Biophys Res Commun* 1996;226:561-5.
15. Criteria for diagnosis of Behçet's disease. International Study Group for Behçet's Disease. *Lancet* 1990;335:1078-80.
16. Akyol O, Canatan H, Yilmaz HR, Yuce H, Ozyurt H, Sogut S, et al. PCR/RFLP-based cost-effective identification of SOD2 signal (leader) sequence polymorphism (Ala-9Val) using NgoM IV: a detailed methodological approach. *Clin Chim Acta* 2004;345:151-9.
17. Mitrunen K, Sillanpää P, Kataja V, Eskelinen M, Kosma VM, Benhamou S, et al. Association between manganese superoxide dismutase (MnSOD) gene polymorphism and breast cancer risk. *Carcinogenesis* 2001;22:827-9.
18. Akyol O, Yanik M, Elyas H, Namli M, Canatan H, Akin H, et al. Association between Ala-9Val polymorphism of Mn-SOD gene and schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2005;29:123-31.
19. Yen JH, Tsai WC, Lin CH, Ou TT, Hu CJ, Liu HW. Cytochrome P450 1A1 and manganese superoxide dismutase gene polymorphisms in Behçet's disease. *J Rheumatol* 2004;31:736-40.
20. Cox DG, Tamimi RM, Hunter DJ. Gene x Gene interaction between MnSOD and GPX-1 and breast cancer risk: a nested case-control study. *BMC Cancer* 2006;6:217.
21. Lightfoot TJ, Skibola CF, Smith AG, Forrest MS, Adamson PJ, Morgan GJ, et al. Polymorphisms in the oxidative stress genes, superoxide dismutase, glutathione peroxidase and catalase and risk of non-Hodgkin's lymphoma. *Haematologica* 2006;91:1222-7.
22. Dedoussis GV, Kanoni S, Panagiotakos DB, Louizou E, Grigoriou E, Chrysohoou C, et al. Age-dependent dichotomous effect of superoxide dismutase Ala16Val polymorphism on oxidized LDL levels. *Exp Mol Med* 2008;40:27-34.
23. Gurel A, Tomac N, Yilmaz HR, Tekedereli I, Akyol O, Armutcu F, et al. The Ala-9Val polymorphism in the mitochondrial targeting sequence (MTS) of the manganese superoxide dismutase gene is not associated with juvenile-onset asthma. *Clin Biochem* 2004;37:1117-20.
24. Aydin E, Sögüt S, Ozyurt H, Ozugurlu F, Akyol O. Comparison of serum nitric oxide, malondialdehyde levels, and antioxidant enzyme activities in Behçet's disease with and without ocular disease. *Ophthalmic Res* 2004;36:177-82.
25. Ilhan A, Iraz M, Gurel A, Armutcu F, Akyol O. Caffeic acid phenethyl ester exerts a neuroprotective effect on CNS against pentylentetrazol-induced seizures in mice. *Neurochem Res* 2004;29:2287-92.
26. Erdogan H, Fadillioglu E, Ozgocmen S, Sogut S, Ozyurt B, Akyol O, et al. Effect of fish oil supplementation on plasma oxidant/antioxidant status in rats. *Prostaglandins Leukot Essent Fatty Acids* 2004;71:149-52.
27. Yildirim Z, Turkoz Y, Kotuk M, Armutcu F, Gurel A, Iraz M, et al. Effects of aminoguanidine and antioxidant erdosteine on bleomycin-induced lung fibrosis in rats. *Nitric Oxide* 2004;11:156-65.
28. Yuce H, Hepsen IF, Tekedereli I, Keskin U, Elyas H, Akyol O. Lack of association between pseudoexfoliation syndrome and manganese superoxide dismutase polymorphism. *Curr Eye Res* 2007;32:387-91.
29. Sutton A, Houry H, Prip-Buus C, Capanec C, Pessayre D, Degoul F. The Ala16Val genetic dimorphism modulates the import of human manganese superoxide dismutase into rat liver mitochondria. *Pharmacogenetics* 2003;13:145-57.