

Antiphospholipid syndrome presenting as chronic venous insufficiency in a young male

Marium Marufia Khan¹ , Amara Zafar¹ , Sana Shahid² 

¹Department of Internal Medicine, Dow University of Health Sciences, Karachi, Pakistan

²Department of General Surgery, Dow University of Health Sciences, Karachi, Pakistan

Antiphospholipid syndrome (APS) is an acquired cause of thrombophilia with persistent elevation of antiphospholipid antibodies which include anticardiolipin antibody and lupus anticoagulant.¹ Vascular thrombosis with obstetric complications is the key feature of APS.² Post-thrombotic syndrome (PTS) presenting as non-healing recurrent ulcers are the most frequent chronic complications of deep venous thrombosis (DVT), accounting for 20 to 40% of patients with DVT.³

Herein, we report a rare presentation of primary APS in a young male as PTS. A 19-year-old male presented with bilateral, non-healing, painless recurrent leg ulcers (Figure 1) and visible abdominal veins (Figure 2) for the last six months. According to the patient, he had recurrent painful swelling of lower limbs from 12 years of age which gradually resolved on some oral medications each time. He was recently diagnosed as hypertensive and was non-compliant to medications. He was a non-smoker. There was no history of prolonged immobilization, recent trauma, or surgery. Family history was unremarkable. On physical examination, all peripheral pulses were

palpable. He had deep and dry ulcers on legs with irregular sloping borders along with surrounding lipodermatosclerotic changes bilaterally and prominent distended abdominal veins in flank region bilaterally with decreased flow. A written informed consent was obtained from the patient.

In the laboratory analysis, hemoglobin level, total leukocyte count, erythrocyte sedimentation rate, C-reactive protein, prothrombin time, and international normalized ratio were all normal. The platelet count was 120,000/mm³ and activated partial thromboplastin time was 66.9 sec. Total proteins, serum albumin, urine analysis, and liver and renal function tests were all normal. Hepatitis (HepA, HepB, HepC) and human immunodeficiency virus (HIV) serology were negative. Lipid profile showed serum cholesterol 189 mg/dL, serum triglyceride 178 mg/dL, high-density lipoprotein cholesterol (HDL) 32 mg/dL, and low-density lipoprotein cholesterol (LDL) 121 mg/dL. Screening for lupus with anti-nuclear antibody (ANA) and extractable nuclear antigen (ENA), and rheumatoid arthritis with rheumatoid factor were both negative. Protein C, Protein S, activated protein C, antithrombin III activity and homocysteine levels

Received: October 22, 2020 **Accepted:** December 28, 2020 **Published online:** January 13, 2021

Correspondence: Amara Zafar, MD. Department of Internal Medicine, Dow University of Health Sciences, 74200 Karachi, Pakistan.
Tel: +92 213-453 38 59 e-mail: amara.zafar@live.com

Citation:

Khan MM, Zafar A, Shahid S. Antiphospholipid syndrome presenting as chronic venous insufficiency in a young male. Arch Rheumatol 2021;36(4):615-616.

©2021 Turkish League Against Rheumatism. All rights reserved.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes (<http://creativecommons.org/licenses/by-nc/4.0/>).



Figure 1. Leg ulcer at presentation.

were within the normal range. However, lupus anticoagulant was 76.1 sec, anticardiolipin antibody immunoglobulin (Ig) G was 63.14 GPL/mL and anticardiolipin antibody IgM was 12.29 MPL/mL. Doppler ultrasound revealed all bilateral leg veins showing partial compressibility with mildly reduced augmentation and flow. Doppler findings of the inferior vena cava, portal vein, hepatic veins, renal arteries, and lower limb arteries were all normal. Abdominal ultrasound revealed normal findings. Doppler examination of the renal arteries was unremarkable. Lupus anticoagulant serology was repeated after 12 weeks which was again found to be positive.

Investigations confirmed the diagnosis of primary APS and excluded other causes of hypercoagulability. Treatment with warfarin 5 mg, subcutaneous enoxaparin 60 mg as the bridging therapy, amlodipine/valsartan 10/160 mg P.O. and atorvastatin 20 mg P.O., all once daily, was initiated. He was advised compression dressing with zinc oxide for venous ulcers.




Figure 2. Distended abdominal veins at presentation.

The patient is still under follow-up on a monthly basis in the outpatient setting. His ulcers have improved significantly, and he is on anticoagulants.

In conclusion, primary APS is a rare disease in adolescents and non-healing recurrent venous ulcers in relation to APS are reported rarely, particularly at a young age in male sex. This case is presented due to its unusual presentation to consider possibility of both acquired and hereditary thrombophilic disorders in any patient presenting with chronic venous insufficiency.

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Mia Rodziewicz, David P D'Cruz, Martin Gulliford, O30 The epidemiology of antiphospholipid syndrome in the UK, 1990-2016. *Rheumatology* 2019;58 Suppl_3:kez105.029.
2. Schreiber K, Sciascia S, de Groot PG, Devreese K, Jacobsen S, Ruiz-Irastorza G, et al. Antiphospholipid syndrome. *Nat Rev Dis Primers* 2018;4:17103.
3. Yamaki T. Post-thrombotic syndrome—Recent aspects of prevention, diagnosis and clinical management. *Reviews in Vascular Medicine* 2016;6:10-9.